

Figure 1.—The distribution of the white-pine cone beetle.

Although the white-pine cone beetle damages mainly the second-year cones, first-year conelets and new shoots are also attacked (fig. 3). The severity of these secondary attacks apparently depends on the number of second-year cones available and the size of the attacking beetle population. At times the beetles will attack the male flowers and the buds.

### Evidence of Infestation

From the middle of May until autumn, the infested cones may be found littering the duff beneath the white pines. These cones are characterized by a small exudation at the junction of the cone and petiole, a mixture of pitch, frass, and cone borings.

The infested second-year cones generally drop to the ground within 6 weeks after attack. However, a heavy rain 1 or 2 weeks after a beetle attack may cause the cones to break away and drop. The break occurs at the junction of the petiole and the twig, presumably as a result of the formation of an abscission layer. Larger cones (those attacked later in the season) break away almost immediately, the break occurring between the cone and petiole.

### Description

The white-pine cone beetle egg is pearl white. Average size is 0.75




F-505546

Figure 2.—Part of a white pine cone, showing typical gallery construction by the white-pine cone beetle adult.

mm. long and 0.05 mm. wide. The larva is legless, soft and fleshy, white or cream colored, and it has a pale amber head. Two larval instars develop, which do not differ greatly except in size. The mean width of head capsule is 0.33 mm. in the first instar and 0.49 mm. in the second.

The pupa is white when first formed. After 4 or 5 days the mouth parts, eye areas, and the ends of the elytra become darkened.

The adult is a small, stout beetle 2.8 mm. to 4.2 mm. long, shiny black, and covered with moderately long, erect hairs. The males and females can be distinguished by



F-505547, 505548

**Figure 3.—A, First-year white pine conelet attacked by white-pine cone beetle; B, cone beetle attacking new growth of white pine.**

differences in the dorsal genital plate. In the male the dorsal plate is split laterally by a suture that divides it into two sclerites. The female genital plate is not so divided. The adult males and females are approximately equal in number.

### Life History and Habits

In the Eastern United States there is one generation of the white-pine cone beetle a year. Seasonal activity begins in late April when the adult beetles emerge from their overwintering habitat and fly to the crowns of nearby white pines. The female beetle initiates the attack on the cone, boring in and constructing a tunnel across the cone axis, thus cutting off the vessels and killing the cone. She then turns and bores along the cone axis, constructing an enlarged chamber just inside the entrance (fig. 2).

The male beetle does not participate in the initial attack and does little to help in gallery construction. He enters the cone either through the female's entrance or by boring in adjacent to it. Within the cone he is usually found behind the female, widening the gallery and packing frass into the proximal chamber.

The type of gallery formed apparently depends on the presence or absence of the male. If no male enters the cone, the female bores out through the side of the cone or leaves through the entrance, to try

again in another cone. But if a male finds and enters the cone, the female stays and extends the gallery the full length of the cone axis and emerges at or near the apex (fig. 4). The male either follows or detours from the main gallery, to bore out through the side of the cone.

Egg-laying usually begins in the second cone attacked. This is 10 to 12 days after the adults emerge from their overwintering cones. Egg niches are constructed along the sides of the gallery. The eggs are deposited singly, and each niche is filled with fresh frass.

The eggs hatch in 7 days. Newly hatched larvae begin feeding on the seed embryo and cone tissue. As



F-505549

Figure 4.—White-pine cone with cone beetle gallery extending full length of the cone axis.

they develop, they continue to devour the cone contents, feeding indiscriminately in no particular pattern (fig. 5). The larval period is about 4 weeks.

Pupation takes place in a small frass-filled chamber at the end of the larval gallery. One week later callow adults may be observed. Immature stages may occur as late as September, but the insects overwinter only as adults.

Some adults may emerge from the brood cones. However, most of the new adults remain in the brood cones to overwinter. The new beetles that do leave the brood cone migrate to the tree crown and attack the current year's conelets or remain on the ground to bore into other infested cones. At this time the males initiate as many attacks as the females.

In the fall, the activity of the beetle differs in attacking conelets. There are two characteristic patterns: (1) The beetle bores distally, stopping just short of the apex, then retreats and tunnels into the petiole; or (2) the beetle bores directly into the petiole. Rarely is more than one beetle found in these conelets.

### Natural Control

The white-pine cone beetle is attacked by a number of parasites and predators. Several lepidopterous insects—including *Dioryctria abietella*, *Eucosma monitorana*,



F-505550

Figure 5.—White-pine cone beetle larva feeding on cone tissue.

*E. tocullionara*, and two species of Blastobasidae—feed in the deteriorating cones and occasionally prey on the cone beetle in its various stages. Hymenopterous eggs, larvae, and pupae have been found in, on, or near cone beetle larvae; but only one species—*Cephalonomia hyalinipennis*—has been reared successfully and identified.

Adult *Spathius* sp. have been observed emerging from cones infested by the cone beetle. The exact relationship of these insects to the cone beetle is not known, but certain species of *Spathius* are parasites of other Scolytidae. Various other beetles and flies have been found in white pine cones. These are mostly scavengers, but some no doubt compete with the cone beetles for space and food.

Nematodes (*Aphelenchoides* sp.) have been found in cone beetles taken from overwintering cones. These have not been determined positively as parasites, but other species of the same genus are reported to be parasitic on Scolytidae.

The failure of a number of cone beetle broods to develop has been attributed to pathogenic organisms. A white fungus, abundant in most cones lying in the duff, may develop so rapidly that the copious mycelium engulfs the cone beetle larva and callow adult, preventing them from feeding and eventually causing their death. *Aspergillus nigrans* has been cultured from tissue extracted from apparently paralyzed cone beetle larvae.

### Chemical Control

Various chemical insecticides have been tried without success against this pest.

### References

ADDITIONS TO THE CHECK-LIST OF THE INSECTS OF CONNECTICUT. W. E. Britton. Conn. State

Geol. and Nat. Hist. Survey Bul. 60, 201 pp. 1938.

STIMULATION OF SEED PRODUCTION OF WHITE PINE (*PINUS STROBUS* L.).\* H. W. Hocker. N.H. Agr. Expt. Sta. Ann. Prog. Rpt., Federal-Grant Proj. NE-27, 1 p. 1961.

A NEW GENUS OF SCOLYTTID BEETLES. A. D. Hopkins. Wash. Acad. Sci. Jour. 5: 429-433. 1915.

DESCRIPTION OF THE PINE-CONE-INHABITING SCOLYTTID. E. A. Schwarz. Ent. Soc. Wash. Proc. 3: 143-145. 1895.

CANADIAN BARK BEETLES, PART II. J. M. Swaine. Canada Dept. Agr. Ent. Branch Bul. 14, 143 pp. 1918.

THE USE OF LARVAL ANATOMY IN THE STUDY OF BARK BEETLES (COLEOPTERA: SCOLYTTIDAE). J. B. Thomas. Canad. Ent. Sup. 5, 45 pp. 1957.

FOREST INSECT AND DISEASE CONDITIONS IN THE NORTHEAST—1956.\* W. E. Waters and Alma M. Waterman. U.S. Forest Serv. Northeast. Forest Expt. Sta., Sta. Paper 94, 23 pp. 1957.

---

\*Address requests for copies to the originating office.

## **Pesticide-Information Disclaimer**

***This page has been added; it is not part  
of the original publication.***

\*\*\*\*\*

This USDA Forest Service *Forest Pest Leaflet* (FPL) or *Forest Insect & Disease Leaflet* (FIDL) - both representing the same publication series - has been reproduced in whole from the original publication as a service of the Montana Department of Natural Resources and Conservation (DNRC) Forest Pest Management program. Both FPLs and FIDLs contain useful and pertinent information on forest insect and disease biology, identification, life cycles, hosts, distribution, and potential management options.

Some FPLs and FIDLs, however, discuss and (or) recommend pesticides that are no longer registered with the U.S. Environmental Protection Agency or are no longer available for use by the general public. Use of these pesticides is neither recommended nor endorsed by the Montana DNRC.

Before using any pesticide be sure to consult either a forest health specialist; state extension agent; your state's Departments of Agriculture, Natural Resources, or Forestry; or other qualified professional or agency with any questions on current pesticide recommendations for forest insects and diseases.